

therapy should be concurrently administered); certain acute and chronic dermatoses of varying etiology to help control associated pruritus, irritation, and inflammation; otitis externa in conjunction with topical medication; allergic states, e.g., hives, urticaria, and insect bites; and shock and shock-like states by intravenous administration.

(iii) *Limitations.* See paragraph (c)(1)(iii) of this section.

(3) *Cats*—(i) *Amount.* 0.03125 to 0.125 milligram daily intravenously, intramuscularly, or subcutaneously.

(ii) *Indications for use.* It is used for the treatment of certain acute and chronic dermatoses of varying etiology to help control associated pruritus, irritation, and inflammation.

(iii) *Limitations.* See paragraph (c)(1)(iii) of this section.

[44 FR 16011, Mar. 16, 1978, as amended at 61 FR 5507, Feb. 13, 1996]

#### § 522.970 Flunixin meglumine solution.

(a) *Specifications.* The drug contains 50 milligrams of flunixin per milliliter of aqueous solution.

(b) *Sponsors.* See 000061 in § 510.600(c) of this chapter for use as in paragraph (d) of this section. See 000856 and 059130 for use as in paragraph (d)(1) of this section only.

(c) *Related tolerances.* See § 556.286 of this chapter.

(d) *Conditions of use*—(1) *Horses*—(i) *Amount.* 0.5 milligram of flunixin per pound of body weight (1 milliliter per 100 pounds) per day.

(ii) *Indications for use.* For alleviation of inflammation and pain associated with musculoskeletal disorders, and alleviation of visceral pain associated with colic.

(iii) *Limitations.* For musculoskeletal disorders, administer intravenously or intramuscularly for up to 5 days. For colic, administer a single dose intravenously—treatment may be repeated when signs of colic recur. Caution: The effect of this drug on pregnancy has not been determined. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Beef cattle and nonlactating dairy cattle*—(i) *Amount.* 1.1 to 2.2 milligrams per kilogram of body weight (0.5 to 1

milligram per pound, 1 to 2 milliliters per 100 pounds), once a day as a single dose or divided into 2 doses administered at 12-hour intervals for up to 3 days.

(ii) *Indications for use.* For control of pyrexia associated with bovine respiratory disease and endotoxemia. Also indicated for control of inflammation in endotoxemia.

(iii) *Limitations.* Do not slaughter for food use within 4 days of last treatment. Not for use in lactating or dry dairy cows. A withdrawal period has not been established for use in preruminating calves. Do not use in calves to be processed for veal. Do not use in bulls intended for breeding as reproductive effects in this class of cattle have not been studied. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[42 FR 39103, Aug. 2, 1977, as amended at 52 FR 7832, Mar. 13, 1987; 60 FR 54942, Oct. 27, 1995; 62 FR 22888, Apr. 28, 1997; 63 FR 38749, July 20, 1998]

#### § 522.995 Fluprostenol sodium injection.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains fluprostenol sodium equivalent to 50 micrograms of fluprostenol.

(b) *Sponsor.* See 000859 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* 0.55 microgram fluprostenol per kilogram of body weight.

(2) *Indications for use.* The drug is used in mares for its luteolytic effect to control the timing of estrus in estrous cycling and in clinically anestrus mares that have a corpus luteum.

(3) *Limitations.* Administer by intramuscular injection only. *Warning:* Not for use in horses intended for food. For veterinary use only. Federal law restricts this drug to use by or on the order of a licensed veterinarian. Women of childbearing age, asthmatics, and persons with bronchial and other respiratory problems should exercise extreme caution when handling this product. In the early stages, women may be unaware of their pregnancies. Fluprostenol is readily absorbed through the skin and can cause abortion and/or bronchospasms. Direct